

Universality in Random Walk Models with Birth and Death

Carl M. Bender

Department of Physics, Washington University, St. Louis, MO 63130

Stefan Boettcher

Department of Physics, Brookhaven National Laboratory, Upton, NY 11973

Peter N. Meisinger

Department of Physics, Washington University, St. Louis, MO 63130

February 1, 2008

Models of random walks are considered in which walkers are born at one location and die at all other locations with uniform death rate. Steady-state distributions of random walkers exhibit dimensionally dependent critical behavior as a function of the birth rate. Exact analytical results for a hyperspherical lattice yield a second-order phase transition with a nontrivial critical exponent for all positive dimensions $D \neq 2, 4$. Numerical studies of hypercubic and fractal lattices indicate that these exact results are universal. Implications for the adsorption transition of polymers at curved interfaces are discussed.

PACS number(s): 05.20.-y, 05.40.+j, 05.50.+q

In this letter we study random walk models in which walkers are born at one location and die everywhere else. We believe that our analytical results for the critical behavior exhibited by *spherically symmetric* random walks are universal and therefore provide insights into the dynamics governing dissipative systems that interact at an interface [1]. As an application, we discuss the adsorption transition of a polymer near an attractive boundary [2].

Our spherically symmetric random walk model uses a lattice that drastically simplifies the treatment of a dissipative system interacting with a curved boundary. For example, we can model polymer growth near a hyperspherical boundary in any dimension $D > 0$ by a random walk on a *one-dimensional* lattice. Such a lattice faithfully represents the spatial entropy of the system. While this lattice is not translationally invariant, it is suitable for the study of spherically symmetric boundary-value problems. To demonstrate the utility of this approach we compare numerical results obtained from hypercubic and fractal lattices with our exact analytical results and show that these results are universal.

A random walk on a lattice is described by a single probability function, $C_{\mathbf{n},t;\mathbf{m}}$, the probability that a random walker who begins at location $\mathbf{m}$ at $t = 0$ will be at location $\mathbf{n}$ at time t . The probability $C_{\mathbf{n},t;\mathbf{m}}$ satisfies a difference equation of generic form:

$$C_{\mathbf{n},t;\mathbf{m}} = \sum_{\{\sigma\}_{\mathbf{n}}} P(\sigma \rightarrow \mathbf{n}) C_{\sigma,t-1;\mathbf{m}}, \quad (1)$$

where $\{\sigma\}_{\mathbf{n}}$ is the set of locations σ adjacent to the location $\mathbf{n}$ and $P(\mathbf{n} \rightarrow \mathbf{n}')$ represents the probability that a walker at the location $\mathbf{n}$ will go to the location $\mathbf{n}'$ in one step. The probability $C_{\mathbf{n},t;\mathbf{m}}$ obeys the initial condition $C_{\mathbf{n},0;\mathbf{m}} = \delta_{\mathbf{n},\mathbf{m}}$. Local conservation of probability is expressed by $\sum_{\{\sigma\}_{\mathbf{n}}} P(\mathbf{n} \rightarrow \sigma) = 1$ for all $\mathbf{n}$.

In this letter we generalize the difference equation (1) to allow for creation and annihilation of random walkers. Specifically, we let random walkers give birth in one location, which we label $\mathbf{0}$, with birth rate a and die in all other locations with uniform death rate z . Walkers are created or destroyed in a given location in proportion to the number of walkers at that site, where a and z are the constants of proportionality. Of course, a acts as a birth rate if $a > 1$; if $a < 1$, it is really a death rate. A similar interpretation applies to z .

Birth rates and death rates are properties of populations rather than of single individuals. Thus, rather than solving an initial-value problem for a single random walker, we study a large population of random walkers. We represent such a population by a distribution $G_{\mathbf{n},t}$, which denotes the number of random walkers at location $\mathbf{n}$ at time t . The distribution $G_{\mathbf{n},t}$ satisfies the same recursion relation as $C_{\mathbf{n},t;\mathbf{m}}$ except for factors of a and z :

$$G_{\mathbf{n},t} = \begin{cases} z \sum_{\{\sigma\}_{\mathbf{n}}} P(\sigma \rightarrow \mathbf{n}) G_{\sigma,t-1} & (\mathbf{n} \notin \{\sigma\}_{\mathbf{0}}), \\ aP(\mathbf{0} \rightarrow \mathbf{n}) G_{\mathbf{0},t-1} + z \sum_{\{\sigma \neq \mathbf{0}\}_{\mathbf{n}}} P(\sigma \rightarrow \mathbf{n}) G_{\sigma,t-1} & (\mathbf{n} \in \{\sigma\}_{\mathbf{0}}). \end{cases} \quad (2)$$

Note that $G_{\mathbf{n},t} \geq 0$ for all $\mathbf{n}$ and t . We are interested in the asymptotic behavior of $G_{\mathbf{n},t}$ as $t \rightarrow \infty$. The specific choice of initial distribution $G_{\mathbf{n},0}$ is unimportant, aside from the requirement that it be normalizable, because a random walk is a diffusive (dissipative) process and details of $G_{\mathbf{n},0}$ are irretrievably lost as time evolves; all initial distributions lead to the same large-time behavior. We are interested in steady-state solutions of Eq. (2); the existence of such solutions imposes a relationship between the birth and death rates.

To describe a dissipative system interacting with a hyperspherical boundary we formulate our random walk model on a spherically symmetric lattice in D -dimensional space [3]. We consider the boundary to lie at the origin. (The general case of a hyperspherical boundary of arbitrary radius is considered in Ref. [10].) We take location n to be the volume bounded by two concentric D -dimensional hyperspherical surfaces of radii R_{n-1} and R_n . If the random walker is at location n at time t , then at time $t+1$ the walker must move outward to location $n+1$ with probability $P_{\text{out}}(n)$ or inward to location $n-1$ with probability $P_{\text{in}}(n)$, where $P_{\text{out}}(n) + P_{\text{in}}(n) = 1$ so that probability is locally conserved. Because a walker at the origin ($n=1$) must move outward, we take $P_{\text{out}}(1) = 1$ and $P_{\text{in}}(1) = 0$. Then, the equation for a D -dimensional random walk in Eq. (2) reduces to a one-dimensional recursion relation:

$$G_{n,t} = \begin{cases} zP_{\text{in}}(n+1)G_{n+1,t-1} + zP_{\text{out}}(n-1)G_{n-1,t-1} & (n \geq 3), \\ zP_{\text{in}}(3)G_{3,t-1} + aG_{n-1,t-1} & (n = 2), \\ zP_{\text{in}}(2)G_{2,t-1} & (n = 1). \end{cases} \quad (3)$$

In Ref. [3] we take the probabilities of moving out or in to be in proportion to the hyperspherical surface areas that can be traversed on each move. Let $S_D(R) = 2\pi^{D/2}R^{D-1}/\Gamma(D/2)$

represent the surface area of a D -dimensional hypersphere. Then, for $n > 1$,

$$P_{\text{out}}(n) = \frac{S_D(R_n)}{S_D(R_n) + S_D(R_{n-1})} \quad \text{and} \quad P_{\text{in}}(n) = \frac{S_D(R_{n-1})}{S_D(R_n) + S_D(R_{n-1})}. \quad (4)$$

For dimensions other than $D = 1$ and $D = 2$, when we take $R_n = n$, the difference equation in (3) cannot be solved in closed form [5]. Thus, we propose that the probabilities in Eqs. (4) be replaced by bilinear functions of n , which are uniformly good approximations to $P_{\text{out}}(n)$ and $P_{\text{in}}(n)$ in the range $D > 0$ when $R_n = n$ [6]:

$$P_{\text{out}}(n) = \frac{n + D - 2}{2n + D - 3} \quad \text{and} \quad P_{\text{in}}(n) = \frac{n - 1}{2n + D - 3}. \quad (5)$$

In Ref. [4] it was shown that this crucial simplification in P_{in} and P_{out} preserves the configurational entropy. Now, Eq. (3) can be solved in closed form for all $D > 0$:

$$G_{n,t} = z^t \frac{\Gamma\left(\frac{D+1}{2}\right) \Gamma(n + D - 2)}{2^{n-2} \Gamma(D) \Gamma\left(n + \frac{D-3}{2}\right)} \oint_C \frac{dy}{2\pi i y} y^{n-t-1} \frac{\left(\frac{z}{a}\right)^{\delta_{1,n}} {}_2F_1\left(\frac{n}{2}, \frac{n+1}{2}; n + \frac{D-1}{2}; y^2\right)}{1 + \left(\frac{z}{a} - 1\right) {}_2F_1\left(\frac{1}{2}, 1; \frac{D+1}{2}; y^2\right)}. \quad (6)$$

From the solution in Eq. (6) we can obtain closed-form expressions for spatial and temporal moments of the random walk.

To study critical behavior in the random walk models in Eq. (2) or Eq. (3) we take F , the fraction of walkers at the boundary $\mathbf{0}$ (the location where random walkers are born), as our order parameter. For the case of polymer adsorption, this observable corresponds to the fraction of the polymer adsorbed on the boundary. To construct the fraction F we let $N_t = \sum_{\mathbf{n}} G_{\mathbf{n},t}$ be the total number of random walkers at time t . We restrict our attention to initial distributions for which N_0 is finite so that N_t is finite for all t . Then, $F_t = G_{\mathbf{0},t}/N_t$ represents the fraction of all random walkers at location $\mathbf{0}$ at time t .

We find that, independent of the choice of $G_{\mathbf{n},0}$, the asymptotic behaviors of N_t and F_t are determined by the values of a and z . Specifically, we obtain the generic (lattice-independent) result that the positive quadrant of the (a, z) plane is partitioned into four distinct regions by two boundary curves as shown in Fig. 1. One boundary curve, labeled B_1 in Fig. 1, is a straight line passing through the origin. To the left of B_1 , F_t vanishes as $t \rightarrow \infty$; to the right of B_1 , F_t approaches a positive finite value as $t \rightarrow \infty$. When the lattice

dimension $D \leq 2$ the equation for the boundary line B_1 is $z = a$; as D increases beyond 2 the boundary line remains straight but the slope of B_1 begins to decrease with increasing D . The transition at $D = 2$ is a reflection of Polya's theorem [7], which states that when $D > 2$ the probability of an individual random walker returning to a location is less than unity.

The second boundary curve shown in Fig. 1 is labeled B_2 . The first part of B_2 is a straight-line segment, $z = 1$, extending from the z axis to the boundary line B_1 . This line segment connects to the second part of B_2 , which is a curve that approaches $z = 0$ as $a \rightarrow \infty$. The equation for this second part depends on D . [For a $D = 1$ lattice, where the probabilities of moving to the left or right are both $\frac{1}{2}$, this curve is given by $z = 2a/(a^2 + 1)$ ($a \geq 1$).] Above B_2 , $N_t \rightarrow \infty$ as $t \rightarrow \infty$; below B_2 , $N_t \rightarrow 0$ as $t \rightarrow \infty$. On B_2 the total number of walkers approaches a finite number $N(a)$ as $t \rightarrow \infty$. On the curved portion of B_2 , $N(a) > 0$; on the straight-line portion of B_2 , $N(a) > 0$ for $D > 2$, while $N(a) = 0$ for $D \leq 2$. This transition at $D = 2$ is yet another manifestation of Polya's theorem.

As we cross the boundary line B_1 , the limiting value of the function F_t as $t \rightarrow \infty$ is continuous. We are particularly interested in crossing from one side of B_1 to the other along the boundary curve B_2 that divides the upper region, where $N_t \rightarrow \infty$, and the lower region, where $N_t \rightarrow 0$ as $t \rightarrow \infty$. The interpretation of $\lim_{t \rightarrow \infty} N_t$ being finite and nonzero is that the distribution $G_{\mathbf{n},t}$ approaches a steady-state, where there is a balance between random walkers being created at location $\mathbf{0}$ and annihilated in all other regions.

We focus on B_2 because it is only on this curve that a steady state is reached as $t \rightarrow \infty$. Along B_2 , $F(a) = \lim_{t \rightarrow \infty} F_t$ undergoes a second-order phase transition at the critical point $(a_c, z_c = 1)$, which is situated at the intersection of B_1 and B_2 . On B_2 , $F(a) = 0$ when $a < a_c$ (even though the limiting value of N_t may be 0), and both $N(a)$ and $F(a)$ are *finite* positive numbers when $a > a_c$. The curved portion of B_2 is the locus of all points in the positive quadrant of the (a, z) plane for which both $N(a)$ and $F(a)$ are finite and nonzero.

Many of the qualitative features of Fig. 1 can be derived directly from an analysis of Eq. (2). To determine the boundary line B_1 we introduce a change of variable: $G_{\mathbf{n},t} = z^t H_{\mathbf{n},t}$.

The distribution $H_{\mathbf{n},t}$ satisfies the recursion relation for a random walk with birth rate a/z at location $\mathbf{0}$ and no births or deaths at any other location. Let Π_0 denote the probability that a random walker at location $\mathbf{0}$ will ever return to location $\mathbf{0}$. Suppose a random walk with birth rate a/z and death rate 1 begins at $t = 0$. Of the $H_{\mathbf{0},0}$ walkers who begin at location $\mathbf{0}$, only a fraction Π_0 of them will return to location $\mathbf{0}$ to give birth to new walkers. Of these new walkers, again only Π_0 of them will return to location $\mathbf{0}$ to give birth again, and so on. Hence, to find the total number of walkers who are ever born we must sum a geometric series whose geometric ratio is the quantity $a\Pi_0/z$. If $a\Pi_0/z < 1$, the geometric series converges and the total number of random walkers ever born is finite. As time t increases, the walkers diffuse away from location $\mathbf{0}$. Thus, as $t \rightarrow \infty$, the ratio

$$F_t = \frac{G_{\mathbf{0},t}}{\sum_{\mathbf{n}} G_{\mathbf{n},t}} = \frac{H_{\mathbf{0},t}}{\sum_{\mathbf{n}} H_{\mathbf{n},t}}$$

vanishes. On the other hand, if $a\Pi_0/z > 1$, both $H_{\mathbf{0},t}$ and $\sum_{\mathbf{n}} H_{\mathbf{n},t}$ diverge at the same rate, and the ratio F_t approaches a nonzero limit (that lies between 0 and 1) as $t \rightarrow \infty$.

The transition between $F_t \rightarrow 0$ and $F_t \rightarrow \text{finite limit}$ occurs on the line $z = a\Pi_0$. This is the equation of the boundary line B_1 . Polya's theorem states that for any random walk $\Pi_0 = 1$ when $D \leq 2$ and $\Pi_0 < 1$ when $D > 2$. This theorem explains the transition in the slope of the line B_1 at $D = 2$ [7]. In particular, since $z_c = 1$ for all $D > 0$, we find $a_c = 1/\Pi_0$.

The shape of the curved part of B_2 depends on the choice of lattice; it is not universal. However, the straight-line portion of B_2 is universal and is easy to understand. Points (a, z) such that $a < a_c$ and z near 1 lie to the left of B_1 . Thus, F_t becomes vanishingly small as $t \rightarrow \infty$. Hence, the effect of the birth rate a on the total number of walkers is negligible. The growth or decay of the total number of walkers depends only on z ; if $z < 1$ then $N_t \rightarrow 0$ as $t \rightarrow \infty$, and if $z > 1$ then $N_t \rightarrow \infty$ as $t \rightarrow \infty$.

On the straight-line portion of the curve B_2 , where $a < a_c$ and $z = 1$, the limiting value of N_t depends on the dimension D . If $D \leq 2$ then $a_c = 1$. Thus, on this portion of B_2 a fraction $1 - a$ of random walkers who arrive at location $\mathbf{0}$ at a given time step must die at the next time step. But by Polya's theorem *all* random walkers visit location $\mathbf{0}$ repeatedly.

Hence, the total number of random walkers N_t must vanish as $t \rightarrow \infty$. On the other hand, if $D > 2$ we have $\Pi_0 < 1$. Thus, the fraction $1 - \Pi_0$ of random walkers who originate in location $\mathbf{0}$ *never return* to $\mathbf{0}$. These random walkers never die because $z = 1$. Hence, N_t approaches a finite positive number as $t \rightarrow \infty$.

There is a change in the form of the transition at $D = 4$. When $D < 4$ the slope of B_2 is continuous. However, when $D > 4$ an elbow appears in B_2 at the critical value a_c . Specifically, when $D > 4$ the slope of B_2 is 0 for $0 \leq a < a_c$; just above a_c the slope abruptly becomes $-\Pi_0/(T - 1)$. Here, T is the expected time for a random walker who begins at location $\mathbf{0}$ to return to $\mathbf{0}$ when $a = 1$ and $z = 1$. A quick heuristic argument yields this result [4]: When $D < 4$ we know that T , the first temporal moment of $C_{\mathbf{0},t,\mathbf{0}}$, is divergent; T is finite when $D > 4$. In a steady state all random walkers at location $\mathbf{0}$ leave this location and in a $z = 1$ model only the fraction Π_0 ever return. The random walkers who return to location $\mathbf{0}$ do so in T steps on average. These returning random walkers experience a death rate z for $T - 1$ of these T steps. Thus, the expected number of random walkers who actually return to location $\mathbf{0}$ is decreased by the factor z^{T-1} . Hence, after T steps we expect to find the number of walkers at location $\mathbf{0}$ to be multiplied by the factor $a\Pi_0 z^{T-1}$. The condition that there be a steady state is therefore $a\Pi_0 z^{T-1} = 1$. Near the critical point $a = a_c + \delta$ and $z = 1 - \epsilon$ as $\delta, \epsilon \rightarrow 0^+$. To first order in δ and ϵ the steady-state condition gives the slope of B_2 as $a \rightarrow a_c^+$. Note that this argument is valid only for $a \geq a_c$.

While the above argument is only valid in the neighborhood of a_c , the same reasoning yields the entire curve $z(a)$ in the limit $D \rightarrow \infty$. In this limit $T = 2$. Hence, the steady-state condition gives $z = a_c/a$.

To determine the nature of the phase transition in $F(a)$ along B_2 , we let $g_{\mathbf{n}} = \lim_{t \rightarrow \infty} G_{\mathbf{n},t}$. The steady-state distribution $g_{\mathbf{n}}$ satisfies the difference equation

$$g_{\mathbf{n}} = \begin{cases} z \sum_{\{\sigma\}_{\mathbf{n}}} P(\sigma \rightarrow \mathbf{n}) g_{\sigma} & (\mathbf{n} \notin \{\sigma\}_{\mathbf{0}}), \\ aP(\mathbf{0} \rightarrow \mathbf{n})g_0 + z \sum_{\{\sigma \neq \mathbf{0}\}_{\mathbf{n}}} P(\sigma \rightarrow \mathbf{n}) g_{\sigma} & (\mathbf{n} \in \{\sigma\}_{\mathbf{0}}), \end{cases} \quad (7)$$

which is the time-independent version of Eq. (2). For a steady-state solution having a finite number of random walkers the sum $\sum_{\mathbf{n}} g_{\mathbf{n}}$ exists. If we sum both sides of Eq. (7) over all

locations, we obtain

$$\sum_{\mathbf{n}} g_{\mathbf{n}} = (a - z)g_0 + z \sum_{\mathbf{n}} g_{\mathbf{n}}. \quad (8)$$

Since the sum $\sum_{\mathbf{n}} g_{\mathbf{n}}$ is nonzero we obtain a formula for $F(a)$ in terms of a and $z(a)$:

$$F(a) = \frac{g_0}{\sum_{\mathbf{n}} g_{\mathbf{n}}} = \frac{1 - z(a)}{a - z(a)}. \quad (9)$$

The result in Eq. (9) is valid on the curved part of B_2 ; it is also valid on the straight-line portion of B_2 when $D > 2$. On B_2 we must treat z as a function of a . We emphasize this dependence by writing $z(a)$ and by treating the fraction F as a function of a only.

We have studied Eq. (7) for several types of lattices. We have determined that along the curve B_2 the steady-state distribution fraction $F(a)$ behaves like

$$F(a) \sim C(D)(a - a_c)^\nu \quad (a \rightarrow a_c^+, D \neq 2, 4). \quad (10)$$

The multiplicative constant $C(D)$ depends on the specific choice of lattice. However, the critical exponent ν is universal and only depends on the dimension D . Using the probabilities in Eqs. (5) for the hyperspherical lattice, we derive in Ref. [4]

$$\nu = \begin{cases} \frac{D}{2-D} & (0 < D < 2), \\ \frac{2}{D-2} & (2 < D < 4), \\ 1 & (D > 4). \end{cases} \quad (11)$$

We have verified the universality of these results numerically for more complicated lattices. We have computed the critical exponent ν for three types of random walks that are not analytically tractable: (i) a spherically symmetric random walk using the probabilities in Eqs. (4) for various values of D ; (ii) a conventional random walk on a $D = 3$ cubic lattice; (iii) a random walk on a recursively defined fractal lattice (Sierpinski carpet) [8] with dimension $D = \frac{\ln 12}{\ln 4}$. The agreement between the predicted value of the critical exponent ν in Eq. (11) and that obtained by computer simulation is excellent, which provides strong evidence that the second-order phase transition at a_c is universal.

There is no critical exponent for the special cases $D = 2, 4$; for the case of a spherically symmetric random walk using the probabilities in Eqs. (5) we find that [4] as $a \rightarrow a_c^+$

$$F(a) \sim \begin{cases} \frac{\text{constant}}{a - a_c} e^{-\frac{2}{a - a_c}} & (D = 2), \\ \frac{a - a_c}{12 \log\left(\frac{1}{a - a_c}\right)} & (D = 4). \end{cases} \quad (12)$$

This work can be applied to the study of polymer growth in the vicinity of an attractive boundary [9]. For a hyperspherical boundary of radius m , this phenomenon is described by a partial difference equation of the same form as that in Eq. (7) [5]. Let $P(\kappa)$ represent the fraction of a polymer that is adsorbed as a function of the attractive boundary potential κ . This fraction is analogous to $F(a)$ in Eq. (9). The potential κ is a monotonic function of the birth rate a , and near the critical point [10]

$$P(\kappa) \propto -\frac{dz(a)}{da} \sim C(D)(a - a_c)^{\nu'}. \quad (13)$$

We find that $\nu' = \nu$ ($D < 2$) and $\nu' = \nu - 1$ ($D > 2$), where ν is given in Eq. (11). Thus, the polymer adsorption fraction $P(\kappa)$ exhibits a first-order transition for $D > 4$ and a tricritical point with logarithmic scaling at $D = 4$. This behavior for $P(\kappa)$ is illustrated in Fig. 2.

The analysis in this paper allows us to make a physical prediction for systems where excluded-volume effects can be neglected: The adsorption fraction for a solution of polymers growing near an approximately spherical attractive boundary (such as a cell membrane) exhibits a second-order phase transition with linear scaling.

Two of us, CMB and PNM, wish to thank the U.S. Department of Energy for financial support under grant number DE-FG02-91-ER40628. SB also thanks the U.S. Department of Energy for support under grant number DE-AC02-76-CH00016.

REFERENCES

- [1] See, for example, G. H. Weiss, *Aspects and Applications of the Random walk*, Chap. 5, (North-Holland, Amsterdam, 1994), and references therein.
- [2] G. H. Weiss and R. J. Rubin, *Adv. Chemical Physics* **52**, 363 (1983); A. Takahashi and M. Kawaguchi, *Adv. Polymer Sci.* **46**, 1 (1982); Yu. S. Lipatov and L. M. Sergeeva, *Adsorption of Polymers*, (Halsted, Jerusalem, 1974); E. A. Dimarzio and M. Bishop, *Biopolymers* **13**, 2331 (1974).
- [3] C. M. Bender, S. Boettcher, and L. R. Mead, *J. Math. Phys.* **35**, 368 (1994) and C. M. Bender, S. Boettcher, and M. Moshe, *J. Math. Phys.* **35**, 4941 (1994).
- [4] C. M. Bender, S. Boettcher, and P. N. Meisinger, submitted to *Phys. Rev. E*.
- [5] S. Boettcher, *Phys. Rev. E* **51**, 3862 (1995); S. Boettcher and M. Moshe, *Phys. Rev. Lett.* **74**, 2410 (1995).
- [6] C. M. Bender, F. Cooper, and P. N. Meisinger, submitted to *Phys. Rev. E*.
- [7] G. Polya, *Math. Ann.* **84**, 149 (1921).
- [8] J. Feder, *Fractals* (Plenum, New York, 1988).
- [9] V. Privman, G. Forgacs, and H. L. Frisch, *Phys. Rev. B* **37**, 9897 (1988), and V. Privman and N. M. Švrakić, *Directed Models of Polymers, Interfaces, and Clusters: Scaling and Finite-Size Properties*, (Springer, Berlin, 1989);
- [10] C. M. Bender, S. Boettcher, and P. N. Meisinger, submitted to *Phys. Rev. E*.

FIGURES

FIG. 1. Generic phase diagram for the (a, z) plane. Shown on the diagram are the boundary curves B_1 and B_2 . To the left of B_1 and on B_1 the fraction of random walkers at location $\mathbf{0}$, F_t , approaches 0 as $t \rightarrow \infty$; to the right of B_1 this fraction approaches a finite positive number as $t \rightarrow \infty$. Above B_2 the total number of random walkers, N_t diverges as $t \rightarrow \infty$; below B_2 the total number of walkers approaches 0 as $t \rightarrow \infty$. On B_2 the distribution of random walkers approaches a steady state as $t \rightarrow \infty$. The critical point $(a_c, z_c = 1)$ lies at the intersection of B_1 and B_2 .

FIG. 2. Plot of the adsorption fraction $P(\kappa)$. For increasing $D < 2$ the scaling exponent increases and the transition becomes weaker until at $D = 2$ exponential scaling is obtained. For increasing $D > 2$ the scaling exponent decreases and the transition becomes stronger again; this is compensated by an increase in the critical binding potential κ that is required to bring about the transition. At $D = 4$ we observe a tricritical point with logarithmic scaling. For $D > 4$ the transition is first order, indicated by a discontinuity (green shaded region) in $P(\kappa)$ across the critical point.



